

Structures of β -Amino Ester Enolates: New Strategies Using the Method of Continuous Variation

Lara R. Liou, Anne J. McNeil, Gilman E. S. Toombes, and David B. Collum*


Department of Chemistry and Chemical Biology, Baker Laboratory, Cornell University, Ithaca, New York 14853-1301

Received March 10, 2008; E-mail: dbc6@cornell.edu

Abstract: The solution structures of four enolates derived from β -amino esters are investigated using ^6Li NMR spectroscopy in conjunction with the method of continuous variation (method of Job). Ensembles of homo- and heteroaggregated enolates are generated by mixing enantiomers of a single enolate (R/S mixtures), opposite antipodes of two different enolates (R/S' mixtures), and the same antipodes of two different enolates (R/R' mixtures). The numbers of observable aggregates and their dependence on the mole fraction of the two enolates confirm the hexamer assignments. Inherent symmetries observable in the ^6Li NMR spectra show the stereochemistry of chelation about the hexagonal drum.

Introduction

Very few reactive intermediates have captivated the attention of organic chemists more than lithium enolates.¹ It is not surprising, therefore, that the underlying physical organic chemistry of enolates has garnered attention as well.^{2,3} Understanding structure–reactivity relationships of lithium enolates, however, requires knowing the structures of the aggregates.^{4–6} From the early structural studies of Jackman⁷ up to the more recent studies of Streitwieser⁸ and others,^{9–11} progress has been made, but it is slow and, at times, contentious.¹¹ The challenge stems in large part from a combination of high aggregate symmetries (cf., 1–4) and Li–O linkages that are opaque to NMR spectroscopy.¹²




As part of a collaboration with Sanofi-Aventis to develop a plant-scale synthesis of Otamixaban,¹³ we were charged


- (1) (a) Green, J. R. *Science of Synthesis*; Georg Thieme Verlag: New York, 2005; Vol. 8a, pp 427–486. (b) Schetter, B.; Mahwald, R. *Angew. Chem., Int. Ed.* **2006**, *45*, 7506. (c) Arya, P.; Qin, H. *Tetrahedron* **2000**, *56*, 917. (d) Caine, D. In *Comprehensive Organic Synthesis*; Trost, B. M., Fleming, I., Eds.; Pergamon: New York, 1989; Vol. 1, p 1. Martin, S. F. In *Comprehensive Organic Synthesis*; Trost, B. M., Fleming, I., Eds.; Pergamon: New York, 1989; Vol. 1, p 475. (e) Plaquevent, J.-C.; Cahard, D.; Guillen, F.; Green, J. R. *Science of Synthesis*; Georg Thieme Verlag: New York, 2005; Vol. 26, pp 463–511. (f) *Comprehensive Organic Functional Group Transformations II*; Katritzky, A. R.; Taylor, R. J. K., Eds.; Elsevier: Oxford, U.K., 1995; pp 834–835. (g) Cativiela, C.; Diaz-de-Villegas, M. D. *Tetrahedron: Asymmetry* **2007**, *18*, 569. (h) Dugger, R. W.; Ragan, J. A.; Ripin, D. H. B. *Org. Process Res. Dev.* **2005**, *9*, 253. (i) Farina, V.; Reeves, J. T.; Senanayake, C. H.; Song, J. J. *Chem. Rev.* **2006**, *106*, 2734. (j) Wu, G.; Huang, M. *Chem. Rev.* **2006**, *106*, 2596.
- (2) For a discussion of prior efforts to elucidate structure–reactivity relationships of enolates and other O–Li species see ref 3.
- (3) Liou, L. R.; McNeil, A. J.; Ramirez, A.; Toombes, G. E. S.; Gruver, J. M.; Collum, D. B. *J. Am. Chem. Soc.* **2008**, *130*, 4859.
- (4) Collum, D. B.; McNeil, A. J.; Ramirez, A. *Angew. Chem., Int. Ed.* **2007**, *49*, 3002.
- (5) (a) Hsieh, H. L.; Quirk, R. P. *Anionic Polymerization: Principles and Practical Applications*; Marcel Dekker: New York, 1996. (b) *Ions and Ion Pairs in Organic Reactions*; Szwarc, M., Ed.; Wiley: New York, 1972; Vols. 1 and 2. (c) Wardell, J. L. In *Comprehensive Organometallic Chemistry*; Wilkinson, G., Stone, F. G. A., Abels, F. W., Eds.; Pergamon: New York, 1982; Vol. 1, Chapter 2. (d) Wakefield, B. J. *The Chemistry of Organolithium Compounds*; Pergamon: New York, 1974. (e) Brown, T. L. *Pure Appl. Chem.* **1970**, *23*, 447.
- (6) The rate law provides the stoichiometry of the transition structure relative to that of the reactants: Edwards, J. O.; Greene, E. F.; Ross, J. J. *Chem. Educ.* **1968**, *45*, 381.

- (7) (a) Jackman, L. M.; Bortiatynski, J. *Adv. Carbanion Chem.* **1992**, *1*, 45. (b) Jackman, L. M.; Lange, B. C. *Tetrahedron* **1977**, *33*, 2737. (c) Jackman, L. M.; Chen, X. *J. Am. Chem. Soc.* **1997**, *119*, 8681. (d) Jackman, L. M.; Petrei, M. M.; Smith, B. D. *J. Am. Chem. Soc.* **1991**, *113*, 3451. (e) Jackman, L. M.; Scarmoutzos, L. M.; DeBrosse, C. W. *J. Am. Chem. Soc.* **1987**, *109*, 5355. (f) Jackman, L. M.; Haddon, R. C. *J. Am. Chem. Soc.* **1973**, *95*, 3687. (g) Jackman, L. M.; DeBrosse, C. W. *J. Am. Chem. Soc.* **1983**, *105*, 4177. (h) Jackman, L. M.; Smith, B. D. *J. Am. Chem. Soc.* **1988**, *110*, 3829. (i) Jackman, L. M.; Rakiewicz, E. F. *J. Am. Chem. Soc.* **1991**, *113*, 1202. (j) Also, see: Quan, W.; Grutzner, J. B. *J. Org. Chem.* **1986**, *51*, 4220.
- (8) (a) Streitwieser, A. *J. Mol. Model.* **2006**, *12*, 673. (b) Streitwieser, A.; Wang, D. Z. *J. Am. Chem. Soc.* **1999**, *121*, 6213. (c) Leung, S. S.-W.; Streitwieser, A. *J. Org. Chem.* **1999**, *64*, 3390. (d) Wang, D. Z.; Kim, Y.-J.; Streitwieser, A. *J. Am. Chem. Soc.* **2000**, *122*, 10754. (e) Kim, Y.-J.; Streitwieser, A. *Org. Lett.* **2002**, *4*, 573. (f) Kim, Y.-J.; Wang, D. Z. *Org. Lett.* **2001**, *3*, 2599. (g) Streitwieser, A.; Leung, S. S.-W.; Kim, Y.-J. *Org. Lett.* **1999**, *1*, 145. (h) Abbotto, A.; Leung, S. S.-W.; Streitwieser, A.; Kilway, K. V. *J. Am. Chem. Soc.* **1998**, *120*, 10807. (i) Leung, S. S.-W.; Streitwieser, A. *J. Am. Chem. Soc.* **1998**, *120*, 10557. (j) Abu-Hasanayn, F.; Streitwieser, A. *J. Org. Chem.* **1998**, *63*, 2954. (k) Abu-Hasanayn, F.; Streitwieser, A. *J. Am. Chem. Soc.* **1996**, *118*, 8136. (l) Gareyev, R.; Ciula, J. C.; Streitwieser, A. *J. Org. Chem.* **1996**, *61*, 4589. (m) Abu-Hasanayn, F.; Stratakis, M.; Streitwieser, A. *J. Org. Chem.* **1995**, *60*, 4688. (n) Dixon, R. E.; Williams, P. G.; Saljoughian, M.; Long, M. A.; Streitwieser, A. *Magn. Reson. Chem.* **1991**, *29*, 509.
- (9) (a) Suzuki, M.; Koyama, H.; Noyori, R. *Bull. Chem. Soc. Jpn.* **2004**, *77*, 259. (b) Suzuki, M.; Koyama, H.; Noyori, R. *Tetrahedron* **2004**, *60*, 1571. (c) Pospisil, P. J.; Wilson, S. R.; Jacobsen, E. N. *J. Am. Chem. Soc.* **1992**, *114*, 7585.
- (10) Biddle, M. M.; Reich, H. J. *J. Org. Chem.* **2006**, *71*, 4031.
- (11) The structure and reactivity of lithium enolates during methacrylate polymerizations have garnered considerable attention and have been reviewed: Zune, C.; Jerome, R. *Prog. Polym. Sci.* **1999**, *24*, 631. Also, see: Baskaran, D. *Prog. Polym. Sci.* **2003**, *28*, 521.

with the task of studying the alkylation of β -amino esters (eq 1).¹⁴




Although the mechanistic insights were interesting, it was the characterization of enolate **5** that proved acutely challenging. We solved the problem by generating an ensemble of aggregates from mixtures of antipodes (*R*)-**5** and (*S*)-**5** (eq 2).



⁶Li NMR spectroscopy showed that the number and relative integrations of the aggregates were highly characteristic of hexamers and inconsistent with monomers, dimers, and tetramers. The protocol formally falls under the rubric of the method of continuous variation¹⁵ (also called the method of Job¹⁶), which has found widespread application in chemistry and biochemistry.¹⁷ This strategy was generalized to a range of enolates representing the aggregates (**2–4**) anticipated from crystallographic studies.^{18,19}


The study described herein focuses on the structures of β -amino ester enolates **6–8**. By analogy to **5**, enolates **6–8** were expected to form hexameric drums of general structure **9**, and indeed, they do. We focused on hexamers because they offer unique opportunities to examine new strategies and tactics.



Background

Applying the method of continuous variation to large ensembles in general and enolates **5–8** in particular presents acute challenges. Merely resolving a large number of species spectroscopically can be problematic given the relatively narrow (~ 3 ppm) spectral range of ⁶Li, constituting a major focus throughout this paper. In addition,

enolates **5–8** present considerable complexities stemming from stereoisomerism within the chelates (cf. **10–12**) and positional isomerism (cf. **13** and **14**) characteristic of the hexagonal drum.²⁰ Indeed, ⁶Li NMR spectra recorded at the lowest attainable NMR probe temperatures (< -100 °C) are often intractably complex. Warming the probe, however, elicits rapid chelate–chelate exchange,²¹ conformational exchange,²² and intraaggregate lithium–lithium exchange, which cause the spectra to simplify.^{23,24} Rapid *intraaggregate* exchange in conjunction with slow *interaggregate* exchange causes each aggregate to appear as only a single ⁶Li resonance and affords separate resonances only for aggregates differing by subunit composition (cf. 4:2 versus 3:3 mixed aggregates, **14** and **15**) or by virtue of their aggregation numbers (tetramers versus hexamers). Monitoring the relative ⁶Li resonance integrations as a function of the mole fraction of the enolate mixtures and subsequent parametric fits^{3,14} afford multicomponent Job plots that prove highly diagnostic of the aggregation state. Representative Job plots and new strategies for exploiting ensembles of β -amino ester enolates are described below.



- (18) (a) Seebach, D. *Angew. Chem., Int. Ed. Engl.* **1988**, *27*, 1624. (b) Setzer, W. N.; Schleyer, P. v. R. *Adv. Organomet. Chem.* **1985**, *24*, 353. (c) Williard, P. G. In *Comprehensive Organic Synthesis*; Trost, B. M., Fleming, I., Eds.; Pergamon: New York, 1991; Vol. 1, Chapter 1.1. (d) Pauer, F.; Power, P. P. In *Lithium Chemistry: A Theoretical and Experimental Overview*; Sapse, A.-M., Schleyer, P. v. R., Eds.; Wiley & Sons: New York, 1995; pp 295–392.
- (19) For crystal structures of hexameric O-lithiated species, see: (a) (Simple enolates) Nichols, M. A.; Leposa, C. M.; Hunter, A. D.; Zeller, M. *J. Chem. Crystallogr.* **2007**, *37*, 825. Rodriques-Delgado, A.; Chen, E. Y.-X. *J. Am. Chem. Soc.* **2005**, *127*, 961. Willard, P. G.; Carpenter, G. B. *J. Am. Chem. Soc.* **1986**, *108*, 462. (b) (Chelating enolates) Chivers, T.; Downard, A.; Yap, G. P. A. *Inorg. Chem.* **1998**, *37*, 5708. Jastrzebski, J. T. B. H.; van Koten, G.; van de Mierop, W. F. *Inorg. Chim. Acta* **1988**, *142*, 169. Maetzke, T.; Hidber, C. P.; Seebach, D. *J. Am. Chem. Soc.* **1990**, *112*, 8248. Waldmueller, D.; Mayer, B.; Braun, M.; Hanuschik, A.; Krueger, C.; Guenot, P. *Chem. Ber.* **1992**, *125*, 2779. (c) (Chelating amino alkoxides) Tian, X.; Woski, M.; Lustig, C.; Pape, T.; Froehlich, R.; Le Van, D.; Bergander, K.; Mitzel, N. W. *Organometallics* **2005**, *24*, 82. Strohmman, C.; Seibel, T.; Schildbach, D. *J. Am. Chem. Soc.* **2004**, *126*, 9876. Armstrong, D. R.; Davies, R. P.; Raithby, P. R.; Snaith, R.; Wheatley, A. E. H. *New J. Chem.* **1999**, *23*, 499. Armstrong, D. R.; Clegg, W.; Hodgson, S. M.; Snaith, R.; Wheatley, A. E. H. *J. Organomet. Chem.* **1998**, *550*, 233. Herberich, G. E.; Spaniol, T. P.; Fischer, A. *Chem. Ber.* **1994**, *127*, 1619. (d) (Phenolates) Rajeswaran, M.; Begley, W. J.; Olson, L. P.; Huo, S. *Polyhedron* **2007**, *26*, 3653. Jackman, L. M.; Cizmeciyan, D.; Williard, P. G.; Nichols, M. A. *J. Am. Chem. Soc.* **1993**, *115*, 6262. (e) (Alkoxides) Nekola, H.; Olbrich, F.; Behrens, U. Z. *Anorg. Allg. Chem.* **2002**, *628*, 2067. Chisholm, M. H.; Drake, S. R.; Naini, A. A.; Streib, W. E. *Polyhedron* **1991**, *10*, 805. (f) (Higher aggregates) Andrews, P. C.; MacLellan, J. G.; Mulvey, R. E.; Nichols, P. J. *J. Chem. Soc., Dalton Trans.* **2002**, *8*, 1651. Maetzke, T.; Seebach, D. *Organometallics* **1990**, *9*, 3032. Allan, J. F.; Nassar, R.; Specht, E.; Beatty, A.; Calin, N.; Henderson, K. W. *J. Am. Chem. Soc.* **2004**, *126*, 484. Vilardo, J. S.; Fanwick, P. E.; Rothwell, I. P. *Polyhedron* **1998**, *17*, 769.

- (12) The rapid relaxation of the highly quadrupolar ¹⁷O nucleus would preclude observing ⁶Li–¹⁷O coupling. For leading references on ⁶Li–¹⁵N and ⁶Li–¹³C see: (a) Günther, H. *J. Braz. Chem. Soc.* **1999**, *10*, 241. (b) Collum, D. B. *Acc. Chem. Res.* **1993**, *26*, 227.
- (13) (a) Guertin, K. R.; et al. *Bioorg. Med. Chem. Lett.* **2002**, *12*, 1671. (b) Czekaj, M.; et al. *Bioorg. Med. Chem. Lett.* **2002**, *12*, 1667. (c) Chandramouli, S. V.; O'Brien, M. K.; Powner, T. H. WO Patent 0040547, 2000. (d) Nagula, G.; Huber, V. J.; Lum, C.; Goodman, B. A. *Org. Lett.* **2000**, *2*, 3527.
- (14) (a) McNeil, A. J.; Toombes, G. E. S.; Gruner, S. M.; Lobkovsky, E.; Collum, D. B.; Chandramouli, S. V.; Vanasse, B. J.; Ayers, T. A. *J. Am. Chem. Soc.* **2004**, *126*, 16559. (b) McNeil, A. J.; Toombes, G. E. S.; Chandramouli, S. V.; Vanasse, B. J.; Ayers, T. A.; O'Brien, M. K.; Lobkovsky, E.; Gruner, S. M.; Marohn, J. A.; Collum, D. B. *J. Am. Chem. Soc.* **2004**, *126*, 5938. (c) McNeil, A. J.; Collum, D. B. *J. Am. Chem. Soc.* **2005**, *127*, 5655.
- (15) Gil, V. M. S.; Oliveira, N. C. *J. Chem. Educ.* **1990**, *67*, 473.
- (16) Job, P. *Ann. Chim.* **1928**, *9*, 113.
- (17) (a) Huang, C. Y. *Methods Enzymol.* **1982**, *87*, 509. (b) Hirose, K. *J. Inclusion Phenom. Macrocyclic Chem.* **2001**, *39*, 193. (c) Likkussar, W.; Boltz, D. F. *Anal. Chem.* **1971**, *43*, 1265.

Table 1. Spectroscopically Distinguishable Aggregates in Binary Mixtures of Enolates^a

<i>R/S</i> mixtures	<i>R/S'</i> mixtures	<i>R/R'</i> mixtures
R₆/S₆	R₆	R₆
R₅S₁/R₁S₅	R₅S'₁	R₅R'₁
R₄S₂/R₂S₄	R₄S'₂	R₄R'₂
R₃S₃	R₃S'₃	R₃R'₃
	R₂S'₄	R₂R'₄
	R₁S'₅	R₁R'₅
	S'₆	R'₆

^aThe *R* and *S* notations refer to enantiomers of a single enolate.²⁶ The prime refers to subunits from a structurally distinct enolate. **R_mS_n**/**R_mS_m** refers to a spectroscopically indistinguishable pair of enantiomers.

Results

We examined three strategies for characterizing enolates 6–8. Table 1 summarizes the predicted number of spectroscopically distinct structural forms observed for hexamers.²⁵ In the first column, we describe mixtures of *R* and *S* enantiomers (so-called *R/S* mixtures) derived from a single β-amino ester enolate. This approach used to characterize enolate 5¹⁴ is extended to enolates 6–8. A stereochemical degeneracy simplifies the ⁶Li NMR spectra. The subsequent two columns tabulate ensembles generated from two structurally distinct enolates derived from opposite stereochemical series (*R/S'* mixtures) and from the same stereochemical series (*R/R'* mixtures). *R/S'* and *R/R'* mixtures lift the stereochemical degeneracy (Table 1). The three strategies present different technical challenges and offer complementary views of the hexameric aggregation state.²⁶

***R/S* Mixtures.** We first examined the structures of enolates 6–8 using *R/S* mixtures derived from a single enolate following the protocol established previously.¹⁴ Enantiomeric pairs of aggregates—**R₆/S₆**, **R₅S₁/R₁S₅**, and **R₄S₂/R₂S₄**—are degenerate, resulting in ensembles comprising four rather than seven observable aggregates (Table 1). (Note: **R_mS_n**/**R_nS_m** refers to

- (20) Positional isomerism does not exist in either cyclic dimers or cubic tetramers.
- (21) For examples of chelate exchange, see: (a) Aubrecht, K. B.; Lucht, B. L.; Collum, D. B. *Organometallics* **1999**, *18*, 2981. (b) Reich, H. J.; Goldenberg, W. S.; Sanders, A. W.; Jantzi, K. L.; Tzschucke, C. C. *J. Am. Chem. Soc.* **2003**, *125*, 3509. (c) Fraenkel, G.; Winchester, W. R. *J. Am. Chem. Soc.* **1989**, *111*, 3794.
- (22) For discussions of conformational equilibrations, see: (a) Clegg, W.; Liddle, S. T.; Snaith, R.; Wheatley, A. E. H. *New J. Chem.* **1998**, *22*, 1323. (b) Boche, G.; Fraenkel, G.; Cabral, J.; Harms, K.; van Eikema Hommes, N. J. R.; Lohrenz, J.; Marsch, M.; Schleyer, P. v. R. *J. Am. Chem. Soc.* **1992**, *114*, 1562. (c) Remenar, J. F.; Lucht, B. L.; Kruglyak, D.; Romesberg, F. E.; Gilchirst, J. H.; Collum, D. B. *J. Org. Chem.* **1997**, *62*, 5748.
- (23) Time averaging of lithium nuclei within aggregates (intraaggregate exchanges) is well precedented: (a) Arvidsson, P. I.; Ahlberg, P.; Hilmersson, G. *Chem.—Eur. J.* **1999**, *5*, 1348. (b) Bauer, W. *J. Am. Chem. Soc.* **1996**, *118*, 5450. (c) Bauer, W.; Griesinger, C. *J. Am. Chem. Soc.* **1993**, *115*, 10871. (d) DeLong, G. T.; Pannell, D. K.; Clarke, M. T.; Thomas, R. D. *J. Am. Chem. Soc.* **1993**, *115*, 7013. (e) Thomas, R. D.; Clarke, M. T.; Jensen, R. M.; Young, T. C. *Organometallics* **1986**, *5*, 1851. (f) Bates, T. F.; Clarke, M. T.; Thomas, R. D. *J. Am. Chem. Soc.* **1988**, *110*, 5109. (g) Fraenkel, G.; Hsu, H.; Su, B. M. In *Lithium: Current Applications in Science, Medicine, and Technology*; Bach, R. O., Ed.; Wiley: New York, 1985; pp 273–289. (h) Heinzer, J.; Oth, J. F. M.; Seebach, D. *Helv. Chim. Acta* **1985**, *68*, 1848. (i) Fraenkel, G.; Henrichs, M.; Hewitt, J. M.; Su, B. M.; Geckle, M. J. *J. Am. Chem. Soc.* **1980**, *102*, 3345. (j) Lucht, B. L.; Collum, D. B. *J. Am. Chem. Soc.* **1996**, *118*, 3529.
- (24) For examples of solvent exchanges, see: (a) Lucht, B. L.; Collum, D. B. *Acc. Chem. Res.* **1999**, *32*, 1035 and references therein.
- (25) Ensembles of dimers and tetramers would afford three and five aggregates, respectively, which would afford two and three spectroscopically distinct resonances, respectively. Dimeric and tetrameric ensembles have been discussed in detail previously.³




Figure 1. Job plot showing relative integrations (I_n)²⁷ of **R_mS_{6-n}** hexamers derived from enolates 5 and 6 as a function of the mole fraction of the *R* enantiomer ($X_{(R),6}$) for enolate 6. The curves correspond to a parametric fit to a hexamer model as described in the text. Samples were 0.10 M total enolate in 9.0 M THF/toluene, and spectra were recorded at -20 °C.





Figure 2. Statistical distribution of *R/S* mixtures of hexameric enolates.

spectroscopically indistinguishable pairs of enantiomerically related aggregates.) Characterization of the isopropyl derivative 6 is illustrated emblematically as follows. ⁶Li NMR spectra recorded on *R/S* mixtures of enolate 6 under conditions optimized for fast intraaggregate exchange and slow interaggregate exchange (9.0 M THF in toluene, -20 °C) reveal four distinct resonances consistent with an ensemble of hexamers and inconsistent with monomers, dimers, and tetramers (Table 1).³ Serial dilution at fixed *R/S* proportions showed that the integrations are concentration independent, confirming a common aggregation number.

The integrations of the resonances corresponding to the **R_mS_n** aggregates of 6 were monitored as a function of the mole fraction of the *R* and *S* antipodes. The equilibria are described

- (26) We refer to the enolates as *R* or *S* according to the structures as follows:



In the case of a phenyl moiety ($R = \text{Ph}$), however, the Cahn–Ingold–Prelog (CIP) rules would formally reverse the stereochemical designation. We have taken the liberty of ignoring the reversal to minimize confusion within the text

- (27) The relative integration (I_n) was previously referred to as the aggregate mole fraction (X_n).¹⁴ We have changed it to avoid using two, distinctly different mole fraction terms and to include provisions for mixtures of aggregates which are not of the same aggregation number.




Figure 3. Job plot showing the relative integrations (I_n)²⁷ of putative R_nS_{4-n} tetramers derived from enolates **5** and **7** as a function of the mole fraction of the *R* enantiomer ($X_{(R)-7}$) for enolate **7**. The curves correspond to a flawed parametric fit to the erroneous tetramer model as described in the text. Samples were 0.10 M total enolate in 9.0 M THF/toluene, and spectra were recorded at -25 °C.




Figure 4. Job plot showing the relative integrations (I_n)²⁷ of R_nS_{6-n} hexamers as a function of the mole fraction of the *R* enantiomer ($X_{(R)-7}$) for enolates **5** and **7**. The curves correspond to a parametric fit to a hexamer model assuming two resonances overlap as labeled and described in the text. Samples were 0.10 M total enolate in 9.0 M THF/toluene, and spectra were recorded at -25 °C.






according to eqs 3–5. A parametric fit affords the Job plot shown in Figure 1. The fit to the hexamer model is excellent. The R_3S_3 form is inordinately stable. Figure 2 displays the curves predicted from a purely statistical model.



The corresponding analysis of benzyl-substituted enolate **7** in 9.0 M THF/toluene at -25 °C revealed only three resonances, consistent with either an ensemble of tetramers (R_4/S_4 , R_3S_1/R_1S_3 , and R_2S_2) or an ensemble of hexamers in which two resonances are unresolved. Failure of the tetramer-based model was suggested by a poor parametric fit (Figure 3). By contrast, the fit to a hexamer model assuming the two resonances

corresponding to the R_3/S_3 and R_4S_2/R_2S_4 hexamers are superimposed afforded a distinct improvement (Figure 4). The latter fit is further justified by the sporadic appearance of the fourth resonance as a shoulder. A similar problem surfaced during the analysis of *R/S* mixtures of phenyl-substituted enolate **8** in which resonances corresponding to R_6 and R_5S_1/R_1S_5 do not resolve in 9.0 M THF/toluene. In the case of **8**, however, all four ^6Li resonances fully resolve in 3.0 M THF/toluene, affording a Job plot of quality comparable to that in Figure 1 consistent with hexamers (Supporting Information).

***R/S'* Mixtures.** We examined mixtures of structurally distinct enolates derived from opposite absolute configuration (*R/S'* mixtures).²⁶ The *R/S'* mixtures should be similar to the *R/S* mixtures except that the enantiomeric degeneracy has been removed: An ensemble of $R_mS'_n$ hexamers would appear as seven discrete aggregates (Table 1). Resolution of seven ^6Li resonances was, at times, more difficult because intraaggregate exchange rates varied slightly among aggregates of different stoichiometries. Parametric fits^{3,14} to the equilibria in eqs 6–10 were carried out in each case. The studies were arbitrarily restricted to mixtures of enolates **6**–**8** with enolate **5**. (No such restriction would exist in the event that an enolate resists characterization.)



Analysis of *R/S'* mixtures of enolates **5** and **6** was reasonably successful. All seven resonances were readily observed. At nearly equimolar concentrations ($X_{(R)-5} = 0.4$ – 0.6), the resonances corresponding to $R_4S'_2$ and $R_2S'_4$ were readily observed but difficult to accurately integrate separately. Accordingly, the two resonances were summed, affording the Job plot illustrated in Figure 5. The increased stability of $R_3S'_3$ was anticipated from results for the *R/S* mixtures (vide supra).

R/S' mixtures derived from enolates **5** and **7** also successfully afforded a Job plot, albeit with the requisite summation of the

(28) For instances in which ensembles were used to investigate homoaggregation, see: (a) Kissling, R. M.; Gagne, M. R. *J. Org. Chem.* **2001**, *66*, 9005. (b) Galiano-Roth, A. S.; Michaelides, E. M.; Collum, D. B. *J. Am. Chem. Soc.* **1988**, *110*, 2658. (c) Reich, H. J.; Goldenberg, W. S.; Gudmundsson, B. Ö.; Sanders, A. W.; Kulicke, K. J.; Simon, K.; Guzei, I. A. *J. Am. Chem. Soc.* **2001**, *123*, 8067. (d) Gilchrist, J. H.; Harrison, A. T.; Fuller, D. J.; Collum, D. B. *J. Am. Chem. Soc.* **1990**, *112*, 4069. (e) Hoffmann, D.; Collum, D. B. *J. Am. Chem. Soc.* **1998**, *120*, 5810. (f) Fraenkel, G.; Henrichs, M.; Hewitt, M.; Su, B. M. *J. Am. Chem. Soc.* **1984**, *106*, 255. (g) Novak, D. P.; Brown, T. L. *J. Am. Chem. Soc.* **1972**, *94*, 3793. (h) Desjardins, S.; Flinois, K.; Oulyadi, H.; Davoust, D.; Giessner-Pretre, C.; Parisel, O.; Maddaluno, J. *Organometallics* **2003**, *22*, 4090. (i) Günther, H. In *Advanced Applications of NMR to Organometallic Chemistry*; Gielen, M., Willem, R., Wrackmeyer, B., Eds.; Wiley & Sons: New York, 1996; pp 247–290. (j) Weingarten, H.; Van Wazer, J. R. *J. Am. Chem. Soc.* **1965**, *87*, 724. (k) Góralski, P.; Legoff, D.; Chabanel, M. *J. Organomet. Chem.* **1993**, *456*, 1.




Figure 5. Job plot showing relative integrations (I_n)²⁸ of $R_n S_{6-n}$ hexamers derived from enolates **5** and **6** as a function of the mole fraction of the *S* enantiomer ($X_{(S)-6}$). The curves correspond to a parametric fit to a hexamer model as described in the text. Samples were 0.10 M total enolate in 3.0 M THF/toluene, and spectra were recorded at -30 °C.




Figure 6. ^6Li NMR spectra recorded on *R/R'* mixtures (the same stereochemical series) derived from enolates **5** and **8**. Samples were 0.10 M total enolate in 3.0 M THF/toluene, and spectra were recorded at -40 °C.

resonances corresponding to $R_4 S'_2$ and $R_2 S'_4$ (Supporting Information). By contrast, *R/S'* mixtures of **5** and **8** afford ensembles of aggregates that are qualitatively familiar, but poor resolution precludes a detailed analysis.

***R/R'* Mixtures.** Mixtures of structurally distinct enolates derived from the same stereochemical series (*R/R'* mixtures) afford an ensemble of seven spectroscopically distinct hexamers. Each of the $R_m R'_n$ hexamers should mimic the optically pure (R_6) forms. Success depends critically on the choice of enolates.

^6Li spectra of *R/R'* mixtures derived from enolates **5** and **8** in the limit of rapid intraaggregate exchange (-40 °C) display seven well-resolved resonances bracketed by the resonances of homo-aggregates of **5** and **8** (Figure 6). The resulting Job plot (Figure 7) is both instructive and aesthetically pleasing. (Figure 7 represents a single parametric fit displayed in two parts for visual clarity.) Superposition of the curves from the parametric fit in Figure 7 with the analogous curves anticipated from a purely statistical distribution shows a striking similarity (Figure 8).




Figure 7. Job plot showing the relative integrations (I_n)²⁸ of $R_n R'_{6-n}$ hexamers of enolates **5** and **8** as a function of the mole fraction of the *R* enantiomer of enolate **8** ($X_{(R)-8}$). The curves correspond to a parametric fit of the data to a hexamer model as described in the text. The single fit is dissected into two separate plots for clarity. Samples were 0.10 M total enolate in 3.0 M THF/toluene, and spectra were recorded at -40 °C.





Figure 8. Superposition of the parametric fit from Figure 7 (red) with the results anticipated from a statistical distribution (black).

R/R' mixtures derived from methyl- and isopropyl-substituted enolates **5** and **6** (respectively) presented minor resolution problems. The ^6Li signals of R_6 and $R_2 R'_1$ ($R = 5$; $R' = 6$) were observable as discrete resonances but were poorly resolved, requiring summation of their integrations when the data were fitted (Supporting Information). Mixtures of the methyl- and benzyl-substituted enolates **5** and **7** afforded a poorly resolved ensemble of aggregates due to a small chemical shift dispersion.

Chart 1^a

^a Li and O have been omitted for clarity, and N = NH₂.

Stereochemistry of Chelation. With the hexameric aggregation state established, it is instructive to examine the ⁶Li NMR spectra recorded in the low-temperature limit in which all intraaggregate exchanges are slow. We discuss the results in the context of the possible structural forms shown in Chart 1. Spectra are archived in the Supporting Information.

Let us first consider the structures of the R_3S_3 hexamer available as the predominant species in equimolar *R/S* mixtures (racemates). R_3S_3 hexamers derived from racemates of **5–8** display inordinate stabilities relative to less symmetric counterparts of unequal stoichiometry. Of the three high-symmetry structural types represented by **A**, **B**, and **C**, isomers **A** and **C** should each display a single ⁶Li resonance, whereas **B** should display a pair of resonances in a 1:1 ratio. Isomer **B** appears to be excluded. In light of the crystal structure of **5** showing a hexamer of type **A**,¹⁴ we deem **A** as the probable form in solution. The orientation of the alkyl substituents away from the perimeter of **A** also logically accounts for the unusual stability.

The $R_3S'_3$ hexamers formed from the *R/S'* mixtures (pseudoracemates) display the same unusual stability as the R_3S_3 aggregates. The reduced symmetry of $R_3S'_3$ hexamers causes the *R* antipodes to reside on one hexagonal face and the *S'* antipodes on the other, resulting in two ⁶Li resonances (1:1). Warming causes the two resonances to coalesce, which we attribute to intraaggregate exchange of the lithium nuclei. Although this experiment does not distinguish **A** and **C**, the consistency with **A** is satisfying.

The homochiral (R_6 and S_6) aggregates appear to be hexagonal drum **D**. The R_6 and S_6 homoaggregates of **6–8** each showed two resonances (1:1) at ≤ -90 °C, consistent with **D** and inconsistent with **E** and **F**. Wrapping the six chelates of **E** in the opposite direction affords a second isomer in which both alkyl groups are in the equatorial positions around the perimeter, but it seems unlikely that the two stereoisomers would be of equal stability. The homoaggregates of **5**, however, initially presented a minor conundrum. Previously reported¹⁴ spectra recorded on solutions of optically pure **5** revealed a single ⁶Li resonance, suggesting that the R_6 and S_6 forms of **5** are *not* type **D**. Results from enolates **6–8** suggested that the rates of chelate exchange may be high for the relatively small methyl-substituted enolate. Indeed, reinvestigation of optically pure enolate **5** showed that the R_6 and S_6 forms display two resonances albeit at < -120 °C and low THF concentration. Therefore, enolates **5–8** are all assigned as hexamer type **D**.

In the case of the *R/S* and *R/S'* mixtures, the R_3S_3 and $R_3S'_3$ are inordinately stable and show distinct preferences for a single positional isomer, resulting in relatively clean spectra with minor mounds in the baseline in equimolar mixtures. By contrast, the *R/R'* mixtures should afford the statistically distributed $R_mR'_n$ aggregates and nearly isoenergetic positional isomers (see **13** and **14**). Each positional isomer could display multiple inequivalent lithium sites (as

many as six per isomer). Indeed, the R/R' mixtures afforded only broad mounds regardless of stoichiometry.

Discussion

We are currently developing methods for characterizing a wide range of lithium enolates, all of which are based on the method of continuous variation.^{3,14} Recent studies, for example, show that standard enolates adorned with further functionality are either dimers or tetramers, depending on the choice coordinating solvent.³ A primary goal of the study described herein was to exploit enolates **5–8** derived from β -amino esters to examine methods and strategies for characterizing highly aggregated enolates. We open the discussion by summarizing the structural assignments. This is followed by a more detailed analysis of *how* ^6Li NMR spectroscopy in conjunction with the method of continuous variation (Job plots) afforded the critical aggregation numbers.

Enolate Structures. Job plots salted throughout the Results and discussed below show that enolates **5–8** are hexameric. We believe that *all* of the homo- and heteroaggregated hexamers are of general structure **9** bearing chelates wrapped around the hexagonal drum in the same direction. Direct support for this assertion is based on ^6Li NMR spectroscopic studies of a smaller subset that includes (1) \mathbf{R}_6 and \mathbf{S}_6 homochiral hexamers derived from optically pure samples (denoted by \mathbf{R} and \mathbf{S}), (2) $\mathbf{R}_3\mathbf{S}_3$ heterochiral hexamers prevalent in racemic samples, (3) $\mathbf{R}_3\mathbf{S}'_3$ hexamers prevalent in equimolar mixtures of structurally distinct enolates of opposite absolute configuration, and (4) all $\mathbf{R}_m\mathbf{R}'_n$ (and $\mathbf{S}_m\mathbf{S}'_n$) hexamers prepared from mixtures of structurally distinct enolates of the same absolute configuration. The structures of the $\mathbf{R}_m\mathbf{S}_n$ and $\mathbf{R}_m\mathbf{S}'_n$ hexamers in which $m \neq n \neq 0$ (e.g., 5:1 and 4:2) are assigned as general structure **9** only by analogy.

One consequence of the hexagonal drum **9** is that subunits of opposite absolute configuration exemplified by $\mathbf{R}_3\mathbf{S}_3$ and $\mathbf{R}_3\mathbf{S}'_3$ mixed hexamers allow optimal disposition of the alkyl side chains as shown in hexamer **A** (Chart 1). Type **A** hexamers are reflected in the X-ray structure of *rac*-**5**¹⁴ and display inordinate stability in solution (*vide infra*). Conversely, the homochiral aggregates \mathbf{R}_6 and \mathbf{S}_6 as well as the $\mathbf{R}_m\mathbf{R}'_n$ hexamers derived from pairs of enolates within the same enantiomeric series appear to be type **D** hexagonal drums in which the β -alkyl substituents point away from the perimeter on one hexagonal face and toward the perimeter on the other.

Method of Job. The most probable forms of lithium enolates—monomers, dimers, tetramers, and hexamers (**1–4**)—are indistinguishable by standard NMR spectroscopic methods due to the high symmetry and opaque Li–O linkages. The aggregate structures emerge when mixtures of structurally related enolates afford ensembles of mixed aggregates. The number of possible aggregates in the ensemble increases with aggregate size and is highly diagnostic (Table 1).²⁸ Thus, the total number of aggregates observed is revealing. The *quantitative* dependence of these ensembles on the mole fraction of the enolate subunits completes a compelling picture.

Chemical Shifts and Resolution. Success of the strategy for determining the aggregation state depends critically on the resolution of the discrete aggregates by ^6Li NMR spectroscopy. Although some problems presented by enolates **5–8** may be specific to the β -amino ester enolates, the lessons are proving useful as the method is extended to other classes of enolates.

Let us first consider the chemical shift dispersion resulting from pairs of enantiomers in R/S mixtures (Figure 9A). A




Figure 9. Chemical shift dispersion of enolate aggregates derived from (A) R/S mixtures, (B) R/S' mixtures, and (C) R/R' mixtures.

relatively small number of resonances (four) arise because there are three degenerate pairs. The ^6Li resonances are bracketed upfield by the homochiral $\mathbf{R}_6/\mathbf{S}_6$ aggregates and downfield by the heterochiral $\mathbf{R}_3\mathbf{S}_3$ aggregate. The chemical shifts of the aggregates reflect the degree of heterochirality. The nature of the hydrocarbon does *not* significantly influence the resolution,²⁹ but the proportion of THF in THF/hydrocarbon mixtures is important. Although one has limited *control* over the resolution, the resolution was adequate in most cases.

R/S' mixtures present a different scenario (Figure 9B). Because the R and S' enantiomers come from structurally different enolates, the degeneracy caused by enantiomerism is eliminated: All seven $\mathbf{R}_m\mathbf{S}'_n$ hexamers are spectroscopically distinct (Table 1). The resonances are bracketed by the homochiral \mathbf{R}_6 and \mathbf{S}'_6 forms upfield and the $\mathbf{R}_3\mathbf{S}'_3$ heterochiral form downfield. Clustering of resonances within the $\mathbf{R}_4\mathbf{S}'_2/\mathbf{R}_2\mathbf{S}'_4$ and $\mathbf{R}_5\mathbf{S}'_1/\mathbf{R}_1\mathbf{S}'_5$ pairs causes some resolution problems that are often alleviated by adjusting the THF concentration. The chemical shift dispersion can be controlled by judicious pairing of enolates based on the chemical shifts of the \mathbf{R}_6 and \mathbf{S}'_6 homoaggregates. At the outset, we had the simple notion that maximally resolved \mathbf{R}_6 and \mathbf{S}'_6 pairs would afford maximally resolved ensembles of mixed aggregates. This assumption was wrong. Widely separated resonances for the \mathbf{R}_6 and \mathbf{S}'_6 cause the resonances of the mixed aggregates to merge unpredictably. The best resolution is observed by pairing \mathbf{R}_6 and \mathbf{S}'_6 aggregates with *minimum* chemical shift separation (as shown in Figure 9B), conditions causing the R/S' mixtures to mimic the R/S mixtures.

The R/R' mixtures afford seven ^6Li resonances bracketed by the resonances of the two homoaggregates (Figure 9C). The chemical shifts of the $\mathbf{R}_m\mathbf{R}'_n$ mixed aggregates correlate predict-

(29) Hydrocarbon effects on the chemical shift, aggregation state, and reactivity of organolithiums can be pronounced. For examples and leading references, see: (a) Godenschwager, P. F.; Collum, D. B. *J. Am. Chem. Soc.* **2007**, *129*, 12023.

ably with the stoichiometry. Resolution is maximized by pairing \mathbf{R}_6 and \mathbf{R}'_6 homoaggregates with the largest chemical shift separation. Tactically, this allows one homoaggregate with an unusual chemical shift—the phenyl-substituted enolate **8**—to be paired with the others. Faced with a particularly difficult resolution problem, one might explicitly design a derivative with a β substituent (such as a naphthyl or phenanthryl group) to impart an especially wide chemical shift dispersion. Additional importance of the R/R' mixtures is underscored by the parametric fits.

Parametric Fits. The Job plots and affiliated parametric fits^{3,14} are fully consistent with hexamers. Although in several instances limited resolution demands that resonances corresponding to two aggregates are summed (see Figures 4 and 5), the underlying math is flexible, and the conclusions appear robust. We remind the reader, however, that the primary goal is to exploit an ensemble of *heteroaggregates* to provide insights into the structures of the *homoaggregates*. To that end, control experiments showing that the proportion of all aggregates is independent of the concentration provide important support to the assignments. The parametric fits resulting from the three protocols offer complementary perspectives of the aggregates.

Using R/S mixtures of a single enolate is the simplest of the three protocols in that only four curves must be fit (Table 1). Enolates **6** and **8** afforded excellent results (Figure 1). Unusual stabilities of the $\mathbf{R}_3\mathbf{S}_3$ aggregates afford distributions that deviate markedly from statistical (Figure 2). Poor resolution for R/S mixtures of enolate **7** afforded a resonance count consistent with tetramers rather than hexamers. Fortunately, the parametric fit to a tetramer model (Figure 3) was decidedly worse than the fit to the hexamer model (Figure 4), and subsequent prodding revealed the missing resonance. Both the number and mole fraction dependencies and absolute concentration independencies of the aggregates are consistent with hexamers and argue strongly that the \mathbf{R}_n and \mathbf{S}_n homoaggregates are necessarily \mathbf{R}_6 and \mathbf{S}_6 .

The R/S' mixtures afford ensembles containing seven potentially distinguishable aggregates (no degenerate pairs). Although we were forced to sum two resonances in some cases, the fits are excellent. One might ask what would happen if an \mathbf{R}_6 hexamer and \mathbf{S}'_4 tetramer were mixed. It seems most probable that an intractable mess would result, but that begs the following question: Would an ensemble containing six hexamers and a tetramer be distinguishable from an ensemble of seven hexamers? Probably not from the parametric fit alone. The independence of the ensemble on the absolute enolate concentration provides compelling support.

Job plots based on R/R' mixtures offer the most convincing view. The seven observable aggregates are nearly statistically distributed type **D** hexamers differing only in the β -alkyl group (Figure 8). This is highly consistent with a type **D** hexamer in which the various alkyl substituents are interchangeable isoenergetically.

Conclusion

The studies described above can be distilled to a single concept: Formation of mixed aggregates from mixtures of

closely related organolithium species provides insight into the spectroscopically opaque homoaggregates.²⁸ The method is an unusually complex application of the method of continuous variation.^{15–17} The strategy relies on high-resolution NMR spectroscopy (⁶Li in this instance) and powerful curve fitting algorithms. We focused on β -aminoester enolates **5–8** because previous studies strongly suggested that they would form hexagonal drums. Such hexamers present challenges that are unique in both detail and severity when compared with dimer- and tetramer-based ensembles.³ Nonetheless, the underlying issues are likely to surface again during studies of other highly functionalized enolates, and strategies developed herein should prove instructive.

Experimental Section

Reagents and Solvents. Substrates were prepared using literature procedures described in the Supporting Information. THF was distilled from solutions containing sodium benzophenone ketyl. Hydrocarbon solvents were distilled from blue solutions containing sodium benzophenone ketyl with approximately 1% tetraglyme to dissolve the ketyl. The diphenylacetic acid used to check *n*-BuLi solution titers was recrystallized from methanol and sublimed at 120 °C under full vacuum.³⁰ [⁶Li]LiHMDS used to prepare the enolates was recrystallized.³¹ Air- and moisture-sensitive materials were manipulated under argon using standard glovebox, vacuum line, and syringe techniques.

Preparation of Compounds. The precursors to **5–7** are literature compounds prepared using an Arndt–Eistert homologation of the corresponding α -amino acids.³² The precursor to **8** was prepared according to a literature procedure.³³ Detailed procedures can be found in the Supporting Information.

Enolization. The enolates were prepared in situ for NMR spectroscopy. After the NMR tube was flame dried, the appropriate stock solutions were added, and the tube was sealed. The samples were warmed to 0 °C for 15–30 min and were thoroughly mixed prior to collection of the spectrum. Integrations were measured using the deconvolution macro included with the VNMR software.

Parametric Fit. Data were fit parametrically to mathematical models described previously.³

Acknowledgment. We thank the National Institutes of Health (GM77167) for direct support of this work as well as Dupont Pharmaceuticals, Merck Research Laboratories, Pfizer, Sanofi-Aventis, R. W. Johnson, Boehringer-Ingelheim, and Schering-Plough for indirect support.

Supporting Information Available: Experimental procedures, NMR spectra, and Job plots. This material is available free of charge via the Internet at <http://pubs.acs.org>.

JA8016957

(30) Kofron, W. G.; Baclawski, L. M. *J. Org. Chem.* **1976**, *41*, 1879.

(31) Romesberg, F. E.; Bernstein, M. P.; Gilchrist, J. H.; Harrison, A. T.; Fuller, D. J.; Collum, D. B. *J. Am. Chem. Soc.* **1993**, *115*, 3475.

(32) Podlech, J.; Seebach, D. *Liebigs Ann.* **1995**, 1217.

(33) (a) Davis, F. A.; Reddy, R. E.; Szewczyk, J. M. *J. Org. Chem.* **1995**, *60*, 7037. (b) Davis, F. A.; Reddy, R. E.; Szewczyk, J. M.; Reddy, G. V.; Portonovo, P. S.; Zhang, H.; Fanelli, D.; Reddy, R. T.; Zhou, P.; Carroll, P. J. *J. Org. Chem.* **1997**, *62*, 2555.